

RESEARCH ARTICLE

First identification of genotypes of *Enterocytozoon bieneusi* (Microsporidia) among symptomatic and asymptomatic children in Mozambique

Aly S. Muadica¹✉, Augusto E. Messa, Jr.²✉, Alejandro Dashti¹✉, Sooria Balasegaram³, Mónica Santin⁴✉, Filomena Manjate², Percina Chirinda², Marcelino Garrine^{2,5}, Delfino Vubil², Sozinho Acácio^{2,6}, Pamela C. Köster¹, Begoña Bailo¹, Tacilta Nhampossa^{2,6}, Rafael Calero-Bernal⁷✉, Jason M. Mwenda⁸, Inácio Mandomando^{2,6*}, David Carmena^{1*}✉



OPEN ACCESS

Citation: Muadica AS, Messa AE, Jr., Dashti A, Balasegaram S, Santin M, Manjate F, et al. (2020) First identification of genotypes of *Enterocytozoon bieneusi* (Microsporidia) among symptomatic and asymptomatic children in Mozambique. PLoS Negl Trop Dis 14(6): e0008419. <https://doi.org/10.1371/journal.pntd.0008419>

Editor: Thuy Le, Duke Global Health Institute, UNITED STATES

Received: January 23, 2020

Accepted: May 23, 2020

Published: June 30, 2020

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pntd.0008419>

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

1 Parasitology Reference and Research Laboratory, National Centre for Microbiology, Health Institute Carlos III, Majadahonda, Madrid, Spain, **2** Centro de Investigação em Saúde de Manhiça (CISM), Maputo, Mozambique, **3** Field Epidemiology Services, National Infection Service, Public Health England, London, United Kingdom, **4** Environmental Microbial and Food Safety Laboratory, Agricultural Research Service, United States Department of Agriculture, Beltsville, Maryland, United States of America, **5** Global Health and Tropical Medicine, Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa (IHMT, UNL), Lisbon, Portugal, **6** Instituto Nacional de Saúde (INS), Ministério da Saúde, Maputo, Mozambique, **7** SALUVET, Department of Animal Health, Faculty of Veterinary, Complutense University of Madrid, Madrid, Spain, **8** Family and Reproductive Health Cluster, World Health Organization (WHO) Regional Office for Africa, Congo Brazaville

✉ These authors contributed equally to this work.

* inacio.mandomando@manhica.net (IM); dacarmena@isciii.es (DC)

Abstract

Enterocytozoon bieneusi is a human pathogen with a broad range of animal hosts. Initially, *E. bieneusi* was considered an emerging opportunistic pathogen in immunocompromised, mainly HIV-infected patients, but it has been increasingly reported in apparently healthy individuals globally. As in other African countries, the molecular epidemiology of *E. bieneusi* in Mozambique remains completely unknown. Therefore, we undertook a study to investigate the occurrence and genetic diversity of *E. bieneusi* infections in children with gastrointestinal symptoms as well as in asymptomatic children in Mozambique. Individual stool specimens were collected from 1,247 children aged between 0 and 14 years-old living in urban and rural settings in Zambézia ($n = 1,097$) and Maputo ($n = 150$) provinces between 2016 and 2019. Samples were analysed for *E. bieneusi* by nested-PCR targeting the internal transcribed spacer (ITS) region of the rRNA gene. All positive amplicons were confirmed and genotyped. Penalised logistic regression (Firth) was used to evaluate risk associations. The overall prevalence of *E. bieneusi* in this children population was 0.7% (9/1,247). A 10-fold higher prevalence was found in Maputo (4.0%; 6/150) than in Zambézia (0.3%; 3/1,097). All *E. bieneusi*-positive samples were from children older than 1-year of age, and most (8/9) from asymptomatic children. Nucleotide sequence analysis of the ITS region revealed the presence of four genotypes, three previously reported (Peru11, $n = 1$; Type IV, $n = 2$, and S2, $n = 2$) and a novel genotype (named HhMzEb1, $n = 4$). Novel genotype HhMzEb1 was

Funding: DC received funding from the Health Institute Carlos III, Ministry of Economy and Competitiveness (Spain), under project PI16CIII/00024. TN received funding from the Fundo Nacional de Investigaç o, Ministry of Health (Mozambique) under project 245-INV. Additional funding was obtained from the USAID Country Office of Mozambique under the Fixed Amount Award Grant No. AID-656-F-16-00002 (IM) and the Centers for Disease Control and Prevention (CDC, USA) through the GAVI Alliance under subcontractor agreement MOA# 870-15 SC (JM). The funders did not play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

identified in both asymptomatic (75%, 3/4) and symptomatic (25%, 1/4) children from a rural area in Maputo province in southern Mozambique. Genotypes HhMzEb1, Peru11, S2, and Type IV belonged to the Group 1 that includes genotypes with low host specificity and the potential for zoonotic and cross-species transmission. Being infected by enteric protozoan parasites and no handwashing were identified as risk associations for *E. bieneusi* infection. This study reports the first investigation of *E. bieneusi* genotypes in Mozambique with the identification of three previously reported genotypes in humans as well as a novel genotype (HhMzEb1). Findings highlight the need to conduct additional research to elucidate the epidemiology of *E. bieneusi* in the country, especially in rural areas where poor hygiene conditions still prevail. Special attention should be paid to the identification of suitable animal and environmental reservoirs of this parasite and to the characterization of transmission pathways.

Author summary

Enterocytozoon bieneusi is an obligate intracellular parasite that infects a wide range of vertebrate hosts. It is the most important etiological agent of human microsporidiasis. Most clinical and epidemiological studies conducted to date have focused on immunodeficient or immunosuppressed individuals including HIV+ patients and solid organ transplant recipients, as in those *E. bieneusi* infection causes life-threatening chronic diarrhoea. In contrast, latent microsporidia infections in immunocompetent individuals have received far less attention. Molecular epidemiological studies in humans and animals have revealed that *E. bieneusi* encompasses a very large diversity of genetic variants (genotypes) with marked differences in host specificity and even geographical distribution. In Mozambique, as in many other African countries, the epidemiology of *E. bieneusi* is completely unknown. Therefore, to identify the occurrence and genetic diversity of this pathogen in Mozambique stool samples were obtained from children, including apparently healthy and symptomatic, in Zamb zia and Maputo provinces and tested for *E. bieneusi* by molecular methods. Results demonstrated the presence of *E. bieneusi* genotypes for the first time in Mozambique. Four genotypes were identified, three genotypes that have previously been reported in humans (Peru11, Type IV, and S2) and a novel genotype (HhMzEb1). Two of the genotypes Peru11, Type IV have also been frequently identified in animals indicating that potentially zoonotic *E. bieneusi* genotypes are inadvertently circulating in the surveyed populations. Additional population genetic studies are needed to elucidate the actual extent of the epidemiology and transmission dynamics of *E. bieneusi* in Mozambique.

Introduction

Microsporidia comprises 200 genera and nearly 1,500 species of spore-forming parasites ubiquitously found in the environment and able to colonize/infect a wide variety of invertebrate and vertebrate hosts [1,2]. Among the 17 Microsporidia species infecting humans *Enterocytozoon bieneusi* is the most frequently reported, causing gastrointestinal infections globally [3]. Infections with *E. bieneusi* in immunocompromised individuals (e.g. patients with AIDS, cancer, organ transplant recipients, and the elderly) are usually associated with chronic diarrhoea,

wasting syndrome, and weight loss. Infections in immunocompetent subjects are often asymptomatic or result in self-limited diarrhoea and malabsorption [4,5]. Extraintestinal disorders and pathologies have also been reported, mainly in lung [6]. In addition, *E. bieneusi* infection in early childhood has been recently linked to impaired growth in children in low-resource settings including African countries such as Malawi, South Africa, or Tanzania [7,8]. *Enterocytozoon bieneusi* has a high genetic diversity and is capable of colonizing/infecting a broad spectrum of mammal and avian species. There are marked differences in host specificities and zoonotic potential among those genotypes [9–11]. Infections are acquired through ingestion of food and water contaminated with *E. bieneusi* spores, or through direct contact with faeces of infected persons and animals or with contaminated soils [2,12,13].

The highly polymorphic ribosomal internal transcribed spacer (ITS) of the ribosomal rRNA (rRNA) gene is the most widely used marker for assessing the genetic diversity within *E. bieneusi* [11]. Based on ITS nucleotide sequences nearly 500 genotypes have been validated according to current nomenclature standards that are distributed into 11 distinct phylogenetic groups [10,14]. Groups 1 and 2 comprise genotypes able to infect a broad range of mammalian species including humans and are, therefore, considered potentially zoonotic. On the other hand, groups 3–11 have strong host specificities and are considered to pose little or null zoonotic risk [11].

The Global Enteric Multicenter Study (GEMS) has provided important information on the aetiology and population-based burden of paediatric diarrheal diseases in sub-Saharan Africa and South Asia. GEMS included in their study enteric protozoan parasites *Giardia duodenalis*, *Cryptosporidium* spp., and *Entamoeba histolytica*, but not the microsporidia *E. bieneusi* [15]. However, in recent years there has been a steady increase in the studies aiming to improve our understanding of the epidemiology of *E. bieneusi* in Africa since this parasite is frequently reported in immunocompromised individuals in African countries (Table 1 and S1 Fig). Importantly, early microscopy-based studies documented *E. bieneusi* prevalences as high as 40–67% in apparently healthy people in Cameroon [16] and Nigeria [17]. These findings indicate that asymptomatic, chronic *E. bieneusi* infections could be more frequent than anticipated. Despite the undoubted progress achieved over the past years, the actual epidemiological situation of *E. bieneusi* in most African countries, including Mozambique, remains largely unknown. To fill this gap, the present study aims to investigate the occurrence and genetic diversity of *E. bieneusi* in symptomatic and asymptomatic paediatric populations in two of the most populous provinces of Central (Zambézia) and Southern (Maputo) Mozambique.

Methods

Ethics statement

Written informed consent was obtained from legal guardians of children voluntarily participating in this survey. All procedures involved in the study were approved by the Ethics Committee of the Health Institute Carlos III under reference number CEI PI 17_2017-v3, and the National Bioethics Committee for Health (CNBS–Comité Nacional de Bioética para Saúde) of Mozambique under reference number 52/CNBS/2017 (participants in Zambézia). Similarly, approval by the Centro de Investigação em Saúde de Manhiça's Institutional Bioethics Committee for Health—CIBS (Ref. CIBS-CISM/075/2015), with final approval by CNBS (Ref. 209/CNBS/15), were obtained for participants in Maputo.

Study design

This study is part of an ongoing research collaborative effort involving the Spanish National Centre for Microbiology (SNCM) in Spain, the Centro de Investigação em Saúde de Manhiça

Table 1. *Enterocytozoon bieneusi* infections reported in humans in African countries including country of origin, type of surveyed population, symptomatology, diagnostic method, and reported prevalence and genotypes. Diagnosis was conducted in stool samples except otherwise indicated.

Country	Surveyed population	Clinical manifestations	Diagnostic method	Prevalence (%)	Genotype(s)	Reference
Cameroon	HIV+/TB+; HIV-/TB+; Immunocompetent	Diarrhoea	LM, IFAT	35.7 (10/28); 24.0 (6/25); 67.5 (85/126)	ND	[16]
	HIV-; HIV+	NS	IFAT, PCR	2.9 (22/758); 0.5 (4/758)	A (8), B (3), CAF4 (5), D (3), Type IV ^c (1)	[36]
	HIV+ (adults)	Diarrhoea, asymptomatic	LM, PCR-RFLP	6.5 (3/46); 4.6 (5/108)	Type IV (4)	[39]
	HIV-	None	PCR	2.6 (5/196)	CAF1 ^d (1), Type IV (4)	[51]
Chad	HIV+ ^a	Diarrhoea	LM, IFAT, PCR	100 (1/1)	ND	[52]
Democratic Republic of the Congo	HIV+ (> 15 yrs.)	Diarrhoea	PCR	7.8 (19/242)	D (1), CAF1 ^d (1), KIN-2 (1), KIN-3 (1), NIA1 (1)	[53]
	AIDS patients (> 15 yrs.)	Diarrhoea	IFAT, PCR	5.1 (9/175)	NIA1 (2)	[54]
	AIDS patients	Diarrhoea, other	LM, IFAT,	2.0 (1/50)	ND	[55]
Democratic Republic of São Tome and Principe	Rural children; Urban inpatient children	Diarrhoea, other	PCR	5.2 (7/134); 8.9 (19/214)	CAF1 ^d (4), KIN-3 (1), Type IV (14), A (2), D (5)	[30]
Ethiopia	HIV+, HIV- (adults)	Diarrhoea	U2B, PCR	14.3 (15/105)	ND	[56]
	HIV+, HIV- (adults)	Diarrhoea, other	LM, PCR	12.3 (30/243)	ND	[57]
Gabon	HIV+ (> 16 yrs.)	Diarrhoea	IFAT, PCR	3.0 (25/822)	A (1), CAF1 (3), CAF2 (1), CAF3 (1), CAF4 (4), D (1), Type IV ^c (4)	[36]
Kenya	HIV+ (adults)	Diarrhoea	U2B	3.0 (1/36)	ND	[58]
Madagascar	Inpatients (adults)	Diarrhoea, asymptomatic	PCR	1.5 (1/67); 2.5 (5/198)	ND	[59]
Malawi	HIV+, HIV-	NS	PCR	100 (37/37) ^b	D (4), Type IV ^c (9), Peru8 (1), S1 (1), S2 (11), S3 (2), S4 (1), S5 (4), S6 (2), UG2145 (1),	[35]
	HIV+ (< 18 yrs.)	Diarrhoea, other	qPCR	37.0 (13/35)	ND	[7]
Mali	HIV+ (adults); HIV- (adults)	Diarrhoea	TEM, LM	32.0 (28/88); 27.0 (3/11)	ND	[60]
	HIV+ Immunocompetent children	Diarrhoea	U2B, IFAT, PCR	14.8 (9/61); 0.0 (0/71)	ND	[61]
Mozambique	Children and adults	Asymptomatic	PCR	9.0 (27/301)	ND	[27]
	Children	Diarrhoea, asymptomatic	PCR	0.3 (1/331); 0.9 (8/916)	HhMzEb1 (4), Peru11 (1), Type IV (2), S2 (2)	This study
Niger	HIV+ (Adults); HIV- (children)	Diarrhoea, other	TEM, LM	7.0 (4/60); 0.8 (8/990)	ND	[62]
	HIV+ (mostly adults)	NS	LM, qPCR	10.5 (24/228)	A (10), CAF1 (2), D (2), E (1), HAN1 (1) Type IV ^c (1), NIA1 (3)	[41]
Nigeria	Rural/urban dwellers	NS	LM	39.6 (80/204)	ND	[17]
	HIV- (children)	Diarrhoea, asymptomatic	PCR	9.3 (4/43)	D (2), Type IV ^c (1), novel K-like (1)	[31]
	HIV+, HIV-	Diarrhoea, other	PCR	6.4 (10/157)	A/Type IV (1), D (3), D/WL7 (1), Type IV (1), WL7 (3), WL7/Type IV (1)	[42]
	HIV+ (adults)	Diarrhoea, other	PCR	26.5 (26/98); 4.3 (52/365)	A (22), CAF2 (2), D (31), D/Type IV (1), EbpA (1), Type IV (14), Nig1 (1), Nig2 (1), Nig3 (1), Nig4 (1), Nig5 (1), Peru8 (1)	[43]
	HIV+	Diarrhoea	PCR	7.7 (10/132)	D (1), Nig2 (2), Peru8 (1), Nig4-like (1), Type IV (5)	[63]

(Continued)

Table 1. (Continued)

Country	Surveyed population	Clinical manifestations	Diagnostic method	Prevalence (%)	Genotype(s)	Reference
	HIV+ (adults on HAART)	Diarrhoea, asymptomatic	PCR	11.1 (12/108); 3.4 (6/177)	Mixed genotypes (3), Nig4 (2), Nig6 (10), Nig7 (2), Type IV (1)	[64]
	HIV+	Diarrhoea, asymptomatic	LM, IFAT, PCR	2.6 (5/193)	B (1), P-like, Peru3, PtEb IV, PtEb V, Type IV, UG2145 (2)	[65]
	HIV+ (adults)	Diarrhoea	PCR	5.5 (5/90)	Type IV (4), unknown mixed infection (1)	[66]
South Africa	HIV+ (inpatients); HIV- (children)	Diarrhoea	LM, PCR-RFLP, qPCR	12.9 (33/255); 4.5 (3/67)	ND	[32]
Tanzania	HIV+ (adults); HIV+, HIV- (children)	Diarrhoea, other; Chronic diarrhoea; Acute diarrhoea; Asymptomatic	LM, TEM	3.5 (3/86); 3.4 (2/59); 0.0 (0/55); 20.0 (4/20)	ND	[34]
Tunisia	Immunocompromised patients	Diarrhoea	LM, PCR	3.5 (3/86)	ND	[67]
	HIV+ (newborn)	Asymptomatic	LM, PCR	100 (1/1)	ND	[68]
	HIV+ (adults); HIV- (adults)	Diarrhoea, other	LM, PCR	20.0 (7/35); 5.3 (3/56)	ND	[69]
	HIV+ (adults)	Diarrhoea	LM, PCR	2.5 (3/119); 5.9 (7/119)	ND	[70]
	HIV+ (mostly adults)	Diarrhoea	LM, PCR	100% (7/7)	B (2), D (4), Peru8 (1)	[71]
	HIV+ (adults)	Diarrhoea, asymptomatic	LM, PCR	19.4 (6/31); 2.8 (2/71)	ND	[72]
Uganda and Zambia	HIV ^{a,b}	Diarrhoea, other	LM, TEM	6.5 (5/77)	ND	[73]
	Children (< 5 yrs)	Diarrhoea, asymptomatic	LM, PCR	17.4 (310/1779); 16.8 (112/667)	Type IV ^c (6), UG2145 (1)	[40]
	HIV+ (children); HIV- (children)	Diarrhoea, other	PCR	76.9 (70/91); 6.6 (10/152)	ND	[33]
Zambia	Rural children	Asymptomatic	U2B, TEM	0.6 (1/176)	ND	[74]
Zimbabwe	HIV+ (adults)	Diarrhoea	LM	10.0 (13/129)	ND	[75]
	HIV+ (adults)	Diarrhoea	LM, PCR	46.0 (34/74)	ND	[76]
	HIV+ (adults)	Diarrhoea	LM, PCR	18.0 (10/55); 51.0 (28/55)	ND	[77]
	HIV-	Diarrhoea, other	LM, PCR	33.0 (2/6)	ND	[78]

HAART, Highly active antiretroviral therapy; HIV, Human immunodeficiency virus; IFAT, immunofluorescence antibody test; LM, light microscopy; ND, not determined; NS, not specified; PCR, polymerase chain reaction; qPCR, real-time PCR; RFLP, restriction fragment length polymorphism; TB, Tuberculosis; TEM, Transmission electron microscopy; U2B, Uvitex 2B fluorescent dye.

^a Patient from Chad diagnosed in France.

^b Diagnosed in an intestinal biopsy.

^c Reported as genotype K.

^d Reported as genotype KIN-1.

<https://doi.org/10.1371/journal.pntd.0008419.t001>

(CISM)/Fundação Manhiça (FM) in Mozambique, and the United States Department of Agriculture to assess the molecular epidemiology of the most frequent enteric protist parasites infecting humans in Mozambique. Two independent studies involving different sampling designs were conducted in Zambézia and Maputo provinces.

Collection of stool samples in Zambézia province

Zambézia (population: 5.1 million; total area: 103,478 km²) is the second most-populous province of Mozambique, located in the central coastal region of the country. A prospective cross-

sectional molecular epidemiological study was carried out between October 2017 and February 2019. All enrolled children were between 3 and 14 years old. A first set of samples were obtained from school children attending 18 public schools with each 35–2,111 (mean: 651) children in 10 of the 22 districts of Zambézia. Voluntary participants were provided with sampling kits (sterile polystyrene plastic flask with spatula and a unique identification number) to obtain individual stool samples. In addition, a second set of samples from children presenting with gastrointestinal complaints (abdominal pain, anal pruritus, bloating, constipation, diarrhoea, flatulence, loss of appetite, nausea, vomiting) seeking medical attention in health care centres ($n = 6$) and hospital settings ($n = 1$) from six districts of the province were collected.

For both sets, an aliquot of each faecal sample was transferred to a TOTAL-FIX stool collection device (Durviz, Valencia, Spain) and shipped to SNCM in Majadahonda (Spain) for downstream molecular analyses.

Collection of stool samples in Maputo province

The study in the Maputo province was conducted in the Manhiça district (population: 0.2 million; total area: 2,373 km²) that is located approximately 80 km north of the capital city Maputo in Southern Mozambique. The Manhiça district has been under a Health and Demographic Surveillance System (HDSS) from CISM since 1996, allowing the link between demographic and clinical data of the population [18].

Stool samples were collected from children under 5 years of age under an ongoing surveillance of diarrheal diseases conducted by the CISM since September 2015 in the Manhiça District Hospital (MDH) and other health peripheral facilities [18,19]. Children under 5 years of age presenting to the peripheral health facilities with moderate-to-severe and less-severe diarrhoea were recruited. Definition of diarrhoeal cases and inclusion criteria to participate in the research study were as previously described [20]. In addition, starting from December 2016, 1 to 3 community controls (asymptomatic children free of diarrhoea episode > 14 days) matched to the index case by age, sex and neighbourhood were identified (using the HDSS databases) and enrolled within 14 days after said index case was enrolled. Stools were collected for both cases and controls and sent to the laboratory for processing. Additionally, stools from children presenting to the sentinel health facility complaining with less-severe diarrhoea and those children presenting to the Xinavane Rural Hospital (XRH), were enrolled starting in April 2017 up to December 2018. Stool samples were collected in sterile flasks, transported to the laboratory, and stored at -80°C without any preservatives for further testing of enteric pathogens as previously described [21].

Data collection

Epidemiological and clinical information was collected in a standardized questionnaire provided as part of the sampling kits and labelled with the same identification number. The questionnaire was completed by researchers interviewing participating children and their caretakers at the time of sampling. The questionnaire included basic demographic characteristics (age, sex, place of living), and potential risk factors including contact with livestock or companion animals, type of drinking water, and information regarding the defecation place. There were some differences between Manhiça and Zambézia questionnaires.

DNA extraction and purification

In stool samples from Zambézia, DNA was isolated from 200 mg of faecal material using the QIAamp Fast DNA stool mini Kit and the QIAcube robot (Qiagen, Hilden, Germany) following the manufacturer's instructions except that treatment of the samples with the lysis buffer

was carried out for 10 minutes at 95°C. Purified DNA samples (200 µL) were stored at -20 °C for downstream molecular analysis. In Manhiça, DNA was manually extracted and purified using the QIAamp DNA Stool Mini Kit from 200 mg of faecal material following the manufacturer's instructions, with the same modification for the lysing step. Purified DNA samples (200 µL) were shipped to the SNCM for molecular testing.

Molecular detection of *Enterocytozoon bieneusi*

To detect *E. bieneusi*, a nested PCR was conducted to amplify a 390-bp fragment including the entire ITS as well as portions of the flanking large and small subunit of the rRNA gene [22]. Primary and secondary reactions (50 µL) were carried out using the outer primer set EBITS3 (5'-GGTCATAGGGATGAAGAG-3') and EBITS4 (5'-TTCGAGTTCTTTCGCGCTC-3'), and the inner primer set EBITS1 (5'-GCTCTGAATATCTATGGCT-3') and EBITS2.4 (5'-ATCGCCGACGGATCCAAGTG-3'), respectively. Reaction mixtures consisted of 1 µL of template DNA, 200 nM of each primer, 2.5 units of MyTAQ DNA polymerase (Bioline GmbH, Luckenwalde, Germany), and 5 µL of MyTAQ Reaction Buffer containing 5 mM dNTPs and 15 mM MgCl₂. Cycling conditions for the primary PCR were as follows: after denaturation at 94°C for 3 min, samples were subjected to 35 cycles of amplification (denaturation at 94°C for 30 s, annealing at 57°C for 30 s, and elongation at 72°C for 40 s), followed by a final extension at 72°C for 10 min. Conditions for the secondary PCR were identical to the primary PCR except only 30 cycles were carried out with an annealing temperature of 55°C. Negative (no DNA template) and positive (*E. bieneusi* PCR-positive samples, genotypes Ebcar2 and EbpA) controls were included in all PCR runs. PCR products were resolved on 2% D5 agarose gels (Conda, Madrid, Spain) stained with Pronasafe (Conda).

Sequence analysis

All amplicons of the expected size were directly sequenced in both directions using the internal primer set in 10 µL reaction mixes using Big Dye chemistries and an ABI 3730xl sequencer analyser (Applied Biosystems, Foster City, CA). Sequencing data were viewed using Chromas Lite version 2.1 software (<https://technelysium.com.au/wp/chromas/>) to generate consensus sequences. The Blast tool (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) was used to compare those sequences with reference sequences deposited at the National Center for Biotechnology Information (NCBI). *Enterocytozoon bieneusi* genotypes were determined using the established nomenclature system based on ITS nucleotide sequence [14]. Sequences obtained in the present study were deposited in the GenBank database under accession numbers MN845065 to MN845068.

Phylogenetic analysis

Nucleotide sequences obtained in this study and *E. bieneusi* nucleotide sequences for all genotypes previously identified in human and animals in Africa as well as appropriate reference sequences to include all *E. bieneusi* groups retrieved from GenBank were aligned with the Clustal W algorithm using MEGA X [23]. Phylogenetic inference was carried out by the Neighbor-Joining (NJ) method as previously described [24], genetic distance was calculated with the Kimura parameter-2 model using MEGA X [23].

Statistical analysis

We analysed the data using EpiData 4.2.0 (EpiData Association, Odense, Denmark) and Stata software, versions 15 (STATA Corp., College Station, Texas, US). We calculated odds ratios

(OR) for associations; a probability (P) value < 0.05 was considered evidence of statistical significance. We examined for possible confounders (change of $> 20\%$ in OR) and interactions. As both populations included selection by symptomatic and asymptomatic populations, we included “a priori” the symptomatic variable in the multivariable model. As the event was rare, and some factors (e.g. *G. duodenalis*) were found in all cases, we used a penalised regression (Firth regression) selecting risk factors with a P -value ≤ 0.2 from the univariable analysis, using Akaike’s information criterion (AIC) and Bayesian information criterion (BIC) to determine selection and evaluate the final model [25].

Results

A total of 1,247 children aged between 0 and 14 years-old were recruited to participate in this study in the Manhiça District in Maputo ($n = 150$) and Zambézia province ($n = 1,097$) in Mozambique. The main socio-demographic and epidemiological features of these paediatric populations are summarized in Table 2.

Table 2. Main socio-demographic and epidemiological features, expressed as frequencies, of the Mozambican children populations ($n = 1,247$) investigated in the present study.

	Zambézia			Manhiça district–Maputo		
	Asymptomatic ($n = 807$)	Symptomatic ($n = 290$)	Total ($n = 1,097$)	Asymptomatic ($n = 109$)	Symptomatic ($n = 41$)	Total ($n = 150$)
Gender						
Male	49.7	50.3	49.9	54.1	48.8	52.7
Female	50.3	49.7	50.1	45.9	51.2	47.3
Age group (years)						
0–5	13.4	33.8	18.8	100	100	100
6–10	64.3	49.3	60.3	NA	NA	NA
11–14	22.3	16.9	20.9	NA	NA	NA
Area						
Rural	86.5	43.8	75.2	100	100	100
Urban	13.5	56.2	24.8	NA	NA	NA
Contact with livestock						
Yes	4.6	14.1	7.1	59 ^b	46.3 ^c	55.3 ^b
No	95.4	85.9	92.9	41 ^b	53.7 ^c	44.7 ^b
Contact with dogs/cats						
Yes	18.8	26.6	20.9	30 ^b	19.5 ^c	26.9 ^b
No	81.2	73.4	79.1	70 ^b	80.5 ^c	73.1 ^b
Main drinking water source						
River/springs	4.3 ^a	4.1	4.3 ^a	2 ^b	2.4	3.6 ^b
Tap	23.2 ^a	54.1	31.4 ^a	91 ^b	80.5	86.5 ^b
Well	72.0 ^a	41.7	64.0 ^a	7 ^b	17.1	9.9 ^b
Defecation place						
Latrine	88.0	76.9	85.1	100 ^c	100 ^d	100 ^e
Outside	12.0	23.1	14.9	0.0 ^c	0.0 ^d	0.0 ^e

NA: not applicable.

^a Information for four children was not available.

^b Information for nine children was not available.

^c Information for one child was not available.

^d Information for five children was not available.

^e Information for six children was not available.

<https://doi.org/10.1371/journal.pntd.0008419.t002>

In Zambézia, 807 asymptomatic schoolchildren and 290 children presenting with gastrointestinal complaints seeking for medical care were investigated in 10 and six districts of the province, respectively (Table 2 and S1 and S2 Tables). The overall male/female ratio was 1.0, and children in the age group 6 to 10 years-old represented 60% of the surveyed individuals. Three out of four children investigated lived in rural areas.

In the rural Manhiça district, 109 asymptomatic children and 41 children presenting with gastrointestinal complaints (11 with moderate-to-severe diarrhoea and 30 with less severe diarrhoea) and seeking for medical care were investigated (Table 2 and S3 and S4 Tables). In this subpopulation, the male/female ratio was 1.1, and children in the age group 12–23 months-old represented 40.7% of the surveyed individuals. Most children (62.7%) were from the area covered by the MDH. All these children lived in rural areas.

Overall, *E. bieneusi* was detected by nested-PCR and confirmed by sequencing in 0.7% (9/1,247, 95% CI: 0.4–1.4) of the investigated children population. The prevalence was more than 10-fold higher in the Manhiça district in Maputo (4.0%, 6/150, 95% CI: 1.8–8.6) compared to all the districts in Zambézia combined (0.3%, 3/1,097, 95% CI: 0.1–0.8). The prevalence for those without symptoms was 0.9% (8/916, 95% CI: 0.4–1.7) and for those with symptoms 0.3% (1/331, 95% CI: 0.04–2.1), and those with diarrhoea 0.4% (1/251, 95% CI: 0.06–2.8). Nucleotide sequences of the ITS revealed the presence of four distinct *E. bieneusi* genotypes, three previously reported in humans (Type IV, S2, and Peru11) and a novel genotype (named HhMzEb1). No mixed infections were observed. Novel genotype HhMzEb1 differed from genotype D (AF101200) by one nucleotide. Genotype S2 nucleotide sequence obtained in this study showed a SNP at the large subunit region when compared with reference sequence (GenBank accession number FJ439678).

The main socio-demographic, epidemiological, and genotypes identified in each of the *E. bieneusi* infections identified in the present study are shown in Table 3. In the Zambézia province, *E. bieneusi* was found in 3 asymptomatic children. Two of them lived in a rural area, attended the same school, but were infected by different genotypes (Peru11 and S2) of the parasite. Both children were co-infected with *G. duodenalis*, *Blastocystis* sp., and *Strongyloides* spp. The third child was infected by the Type IV genotype of *E. bieneusi* and was co-infected with *G. duodenalis*. None of the symptomatic children investigated in this province tested positive to *E. bieneusi*.

In Maputo province, *E. bieneusi* was found in six male children, one presenting with moderate-to-severe diarrhoea and 5 asymptomatic. Five out of the six cases were younger than 2 years old. The S2 and Type IV genotypes were found in a single boy each, one of them co-infected with Rotavirus. Among the 4 boys infected with the novel genotype, HhMzEb1, one was HIV positive, presented with moderate-to-severe diarrhoea, and was co-infected with *Cryptosporidium meleagridis*. Another boy with this genotype was co-infected with *C. parvum*, and a third one was co-infected with Rotavirus. All six *E. bieneusi*-positive samples were co-infected with *G. duodenalis*.

Univariable analysis (Table 4) of the combined data showed that significant risk associations included infections with *G. duodenalis* or *Cryptosporidium* spp. and age, but handwashing was protective. In the Zambézia dataset alone, *G. duodenalis* was a significant association. In addition, travel, *Blastocystis* sp., and absence of a latrine were associated with presence of *E. bieneusi* (P value < 0.5), but confidence intervals crossed 1, with only 3 positive cases. In the Maputo dataset there were no significant associations. In the final multivariable model using the combined dataset, after adjusting for symptoms, the risk associations for *E. bieneusi* were infection with *G. duodenalis* [OR: 19.11, P = 0.43, 95% CI (1.10–332.94)]; infection with *Cryptosporidium* spp. [OR 5.75, P = 0.49 95% CI (1.01–32.89)] and handwashing [OR 0.31 P = 0.128 95% CI (0.67–1.41)]. Although handwashing was not significant, it was retained in

Table 3. Main socio-demographic features, risk factors, and genotyping data of Mozambican children positive to *Enterocytozoon bieneusi* (n = 9) in the present study.

Province	District	Sample ID	Area	Gender	Age	Contact with livestock and/or poultry	Contact with companion animals	Main source of drinking water	Defecation place	Genotype	GenBank accession number
Zambézia	Gurúe	18	Urban	Male	132 months-old	No	Cat, dog	Public tap	Latrine	Type IV	MN845065
	Mocuba	207	Rural	Male	60 months-old	No	No	Well	Outside	Peru11	MN845067
	Mocuba	210	Rural	Female	60 months-old	No	No	Well	Outside	S2	MN845066
Maputo	Manhiça-MDH	1728661.9 ^{a, c, d}	Rural	Male	16 months-old	Poultry	No	Public tap	Latrine	HhMzEb1 ^b	MN845068
	Manhiça-MDH	1725028.3	Rural	Male	28 months-old	ND	ND	ND	Latrine	HhMzEb1 ^b	Identical to MN845068
	Manhiça- XRH	1754917.2 ^c	Rural	Male	16 months-old	ND	ND	ND	Latrine	HhMzEb1 ^b	Identical to MN845068
	Manhiça- XRH	1725031.3 ^{d, e}	Rural	Male	12 months-old	No	No	Public tap	Latrine	Type IV	Identical to MN845065
	Manhiça- XRH	1754898.4	Rural	Male	12 months-old	Goat, poultry	No	Unprotected well	Latrine	S2	Identical to MN845066
	Manhiça-MDH	1754369.9 ^f	Rural	Male	15 months-old	No	No	ND	Latrine	HhMzEb1 ^b	Identical to MN845068

ND: no data available; MDH: Manhiça District Hospital; XRH: Xinavane Rural Hospital.

^a Symptomatic child presenting with moderate-to-severe diarrhoea.

^b Novel genotype.

^c Co-infected with *Cryptosporidium meleagridis* and HIV positive.

^d Deceased.

^e Co-infected with Rotavirus.

^f Co-infected with *Cryptosporidium parvum*.

<https://doi.org/10.1371/journal.pntd.0008419.t003>

the final model as it confounded the relationship with *Cryptosporidium* spp. Age was not a factor after adjustment for other variables.

Phylogenetic analysis revealed that the novel genotype HhMzEB1 clustered within Group 1 when its relationship with other genotypes of human and animal origin previously reported in Africa was evaluated (Fig 1).

Discussion

Little information is currently available on the occurrence and distribution of enteric parasites in Mozambique [26–28], although the GEMS project has significantly improved our understanding of the epidemiology of diarrhoea-associated with parasites *G. duodenalis*, *Cryptosporidium* spp., and *E. histolytica* [15, 29]. However, there is almost no information regarding Microsporidia, and only one study that aimed to investigate the prevalence of intestinal parasite infections in an informal settlement in Beira, Mozambique has reported *E. bieneusi* [27].

Table 4. Descriptive and univariable analysis of the variables of interest potentially associated with an increased exposure risk to *Enterocytozoon bieneusi* in the present study.

Variable	Cases of <i>E. bieneusi</i> with variable	%	Non cases with variable	%	Odds ratio	95% CI	P value
<i>Cryptosporidium</i> spp.	2	22	35	3	9.82	0.96–53.84	0.001
<i>Giardia duodenalis</i>	9	100	556	45	15.27	2.87–	0.001
Age	Continuous				0.66	0.51–0.85	0.001
Handwashing (yes)	6	67	1152	93	0.14	0.03–0.86	0.001
Male Sex	7	78	622	50	3.47	0.66–34.29	0.1
Has livestock	2	22	150	12	3.61	0.32–25.37	0.115
Water source (not tap water)	3	33	885	71	0.39	0.05–2.94	0.236
Symptomatic	1	11	330	27	0.34	0.01–2.58	0.293
<i>Blastocystis</i> sp.	2	22	156	13	1.98	0.20–10.51	0.387
No latrine	2	22	162	13	1.89	0.19–10.01	0.423
Rural	8	89	967	78	2.24	0.30–99.83	0.435
Has pets (dog/cats)	1	11	267	22	0.72	0.02–6.50	0.766
<i>Strongyloides stercoralis</i>	2	22	253	20	1.11	0.11–5.88	0.895
<i>Entamoeba dispar</i>	1	11	122	10	1.14	0.03–8.65	0.9

95% CI: 95% Confidence Interval.

<https://doi.org/10.1371/journal.pntd.0008419.t004>

They detected *E. bieneusi* by real-time PCR in 9.3% (28/301) of the individuals tested and no genotyping was provided. Therefore, this is to our knowledge the first molecular epidemiological study describing occurrence and genetic diversity of *E. bieneusi* in Mozambique. The overall prevalence of *E. bieneusi* infection in the paediatric (0–14 years of age) populations from Mozambique tested in this study was 0.7% (9/1,247). Infections were more commonly identified in asymptomatic children in rural settings.

The *E. bieneusi* infection rates found in our paediatric populations in Zambézia (0.3%) and Maputo (4%) were lower than the prevalence reported in Beira in Sofala province (9%) [27], indicating potential differences in the geographical distribution of the parasite in Mozambique. The disparity could also be associated with differences in socioeconomic characteristics of the studied population as the study in Beira was conducted in an informal settlement in an area that is frequently flooded and not connected to a sewage system after a local hospital noticed high number of diarrhoea cases in this settlement. Prevalence in Mozambique (0.7%) is lower to those previously documented in immunocompetent children populations (range: 4–9%) in other African countries including the Democratic Republic of São Tomé and Príncipe [30], Nigeria [31], South Africa [32], and Uganda [33], and much lower than those found in similar paediatric populations (range: 20–67%) in Cameroon [16], Nigeria [17], and Tanzania [34] (see Table 1). However, prevalence is influenced by the age group selected in the study and the presence of symptoms, so comparisons among studies must keep those differences in mind.

It is noteworthy to indicate that all *E. bieneusi*-positive children were also co-infected by other pathogens including two children in Manhiça that were HIV-positive. Indeed, one to three additional enteric parasites (*G. duodenalis*, *Cryptosporidium* spp., *Blastocystis* sp., and *Strongyloides* spp.) were found and the strongest risk associations were with *G. duodenalis* and *Cryptosporidium* spp. These data clearly depict a highly endemic scenario where polyparasitism is common as previously reported in the city of Beira [27].

Our study is limited by the small number of positive samples, thus the wide confidence intervals for our results. Furthermore, detailed information on handwashing (use of soap or not) and latrine use (toilet or outside latrine) were unavailable in the Zambézia dataset, and




Fig 1. Phylogenetic relationships among *Enterocytozoon bieneusi* genotypes identified in this study. All genotypes identified in humans and animals in Africa, and genotypes to cover all groups of *E. bieneusi* were included for comparative purposes. Analyses were inferred by a Neighbor-Joining method of the entire ITS of rRNA gene based on genetic distances calculated by the Kimura two-parameter model (MEGA X software). All nucleotide sequences include host information with the GenBank accession number in parenthesis. Nucleotide sequences determined in this study are identified with black circles before the genotype name. White circles indicate genotypes identified in Africa.

<https://doi.org/10.1371/journal.pntd.0008419.g001>

missing values for water source and animal livestock in the Maputo dataset implied limited exploration of these associations, especially given the low numbers of cases.

Nucleotide sequence analysis on the ITS region of the nine positive cases revealed the presence of four genotypes, three previously reported (Peru11, Type IV, and S2) and a novel genotype (named HhMzEb1). This constitutes the first report of these genotypes in Mozambique. Genotypes Type IV and S2 were both identified in Zambézia and Maputo, while Peru11 was only identified in Zambézia and novel genotype HhMzEb1 only in Maputo. All genotypes identified in the present study, including novel HhMzEb1, belonged to Group 1 (Fig 1) that includes *E. bieneusi* genotypes with low host specificity that are found not only in humans but also in domestic and wild animals worldwide [10,11,14]. This is the second report of genotype S2 in humans, that had only been previously reported in Malawi from three children (2 HIV-positive and one HIV-negative) and eight HIV-positive adults [35]. Genotype Type IV (also reported previously as K, Peru2, BEB5, CMITS1, BEB-var, PtEB III) has widely been found in humans in different regions throughout the world [5,36–38], with multiple reports in African countries that include Gabon [36], Cameroon [36,39], Uganda [40], Niger [41], Nigeria [31,42,43], Democratic Republic of São Tomé and Príncipe [30], and Malawi [35] (S5 Table). In addition, Type IV has also been reported in a wide range of animal hosts including non-human primates, cattle, cats, dogs, rodents, birds, snakes, and black bears (S5 Table) as well as in water samples [44–47]. There are no previous reports of Peru11 (also reported previously as Peru12) in humans in Africa, but it has been found in humans in Asia and South America (S5 Table). In Africa, Peru11 has been found in a baboon in Kenya [48]. In addition, there are also reports of this genotype in other parts of the world in non-human primates, cats, raccoons, rabbits, rodents, and birds (S5 Table).

In this study, no mixed infections by different *E. bieneusi* genotypes were noticed using ITS-PCR and Sanger sequencing. However, it should be noted that co-infections seem a common finding in human populations in endemic areas. Indeed, intra-isolate diversity has been demonstrated in children with diarrhoea in Uganda using microsatellite markers (in addition to ITS) and subsequent cloning and sequencing of PCR products [49]. These data preclude us to conclusively state that mixed infections are not present in the surveyed Mozambican population as the methodology used in this study was not chosen to address mixed infections. The identification of potentially zoonotic genotypes Peru11 and Type IV, commonly found not only in humans but also in domestic and wild animals worldwide, indicates the potential for zoonotic or cross-species transmission. There are few studies in the African continent that include molecular characterization of *E. bieneusi* in humans and in animals. However, environmental (water) samples should be also considered as a source of infection by microsporidia, as *E. bieneusi* has been demonstrated to be involved in waterborne outbreaks of gastrointestinal disease [50]. Our findings emphasize the need of further studies to explore risk associations as these were limited in our study by the low prevalence and missing information. Studies could include water as well as animals that may be in contact with human populations in Mozambique to understand the modes of transmission of *E. bieneusi*, and, by analogy, of other diarrhoea-causing enteric protist species.

Conclusions

This is the first molecular study of *E. bieneusi* in Mozambique. The parasite was primarily found in asymptomatic children in Maputo (Manhiça district) and Zambézia province. Molecular characterization detected a novel genotype (HhMzEb1) and three known genotypes (Type IV, Peru11 and S2) in diarrheal and healthy Mozambican children. The identification of genotypes previously described in animals suggests potential zoonotic and anthroponotic transmission. There is a lack of information of *E. bieneusi* in the African continent in humans, animal hosts and environmental samples. Under the One Health perspective, further genotyping studies are needed to better understand the epidemiology of this parasite.

Supporting information

S1 Fig. Occurrence of *Enterocytozoon bieneusi* in humans in Africa. Average values for symptomatic and asymptomatic individuals are represented according to reported infection rates summarized in [Table 1](#).

(TIF)

S1 Table. Main socio-demographic features and risk factors of the asymptomatic schoolchildren population ($n = 807$) investigated in Zambézia province (Mozambique), 2017–2019.

(DOCX)

S2 Table. Main socio-demographic features and risk factors of the symptomatic schoolchildren population ($n = 290$) attended at public health centres in Zambézia province (Mozambique), 2017–2018.

(DOCX)

S3 Table. Main socio-demographic features and risk factors of the asymptomatic children population ($n = 109$) investigated in Maputo province (Mozambique), 2016–2018.

(DOCX)

S4 Table. Main socio-demographic features and risk factors of the symptomatic children population ($n = 41$) investigated in Maputo province (Mozambique), 2016–2018.

(DOCX)

S5 Table. Summary of all *Enterocytozoon bieneusi* genotypes reported in Africa including host and geographic range. Highlighted in bold are African countries.

(DOCX)

Acknowledgments

We thank the children and their caretakers who participated in the study, as well as the clinical, field and laboratory staff who worked tirelessly to ensure the data collection and laboratory testing was performed according to the protocols. We also thank all the local government authorities (district Administration Health and Education Directorates in Zambézia and Manhiça) and all community leaders for supporting the study.

Author Contributions

Conceptualization: Inácio Mandomando, David Carmena.

Data curation: Inácio Mandomando, David Carmena.

Formal analysis: Aly S. Muadica, Augusto E. Messa, Jr., Alejandro Dashti, Sooria Balasegaram, Mónica Santin, Rafael Calero-Bernal.

Funding acquisition: Tacilta Nhampossa, Jason M. Mwenda, Inácio Mandomando, David Carmena.

Investigation: Aly S. Muadica, Augusto E. Messa, Jr., Alejandro Dashti, Filomena Manjate, Percina Chirinda, Marcelino Garrine, Delfino Vubil, Sozinho Acácio, Pamela C. Köster, Begoña Bailo, Tacilta Nhampossa.

Methodology: Sooria Balasegaram, Mónica Santin, Rafael Calero-Bernal, Inácio Mandomando.

Project administration: Percina Chirinda, Tacilta Nhampossa, Inácio Mandomando, David Carmena.

Supervision: Marcelino Garrine, Delfino Vubil, Sozinho Acácio, Tacilta Nhampossa, Inácio Mandomando, David Carmena.

Validation: Sooria Balasegaram, Mónica Santin, Rafael Calero-Bernal.

Visualization: Mónica Santin, Rafael Calero-Bernal, Inácio Mandomando, David Carmena.

Writing – original draft: Aly S. Muadica, Augusto E. Messa, Jr., Mónica Santin, Rafael Calero-Bernal, Jason M. Mwenda, Inácio Mandomando, David Carmena.

Writing – review & editing: Aly S. Muadica, Augusto E. Messa, Jr., Sooria Balasegaram, Mónica Santin, Rafael Calero-Bernal, Jason M. Mwenda, Inácio Mandomando, David Carmena.

References

1. Didier ES, Weiss LM. Microsporidiosis: current status. *Curr Opin Infect Dis.* 2006; 19: 485–492. <https://doi.org/10.1097/01.qco.0000244055.46382.23> PMID: 16940873
2. Stentiford GD, Becnel JJ, Weiss LM, Keeling PJ, Didier ES, Williams BAP, et al. Microsporidia—emerging pathogens in the global food chain. *Trends Parasitol.* 2016; 32: 336–348. <https://doi.org/10.1016/j.pt.2015.12.004> PMID: 26796229
3. Weiss LM, Becnel JL. *Microsporidia: Pathogens of Opportunity.* New Jersey: Willey-Blackwell; 2014.
4. Lobo ML, Xiao L, Antunes F, Matos O. Microsporidia as emerging pathogens and the implication for public health: a 10-year study on HIV-positive and negative patients. *Int J Parasitol.* 2012; 42: 197–205. <https://doi.org/10.1016/j.ijpara.2011.12.002> PMID: 22265899
5. Santin M. *Enterocytozoon bieneusi.* In: Xiao L, Ryan U, Feng Y, editors. *Biology of foodborne parasites.* Boca Raton: CRC Press; 2015. pp 149–174.
6. del Águila C, Lopez-Velez R, Fenoy S, Turrientes C, Cobo J, Navajas R, et al. Identification of *Enterocytozoon bieneusi* spores in respiratory samples from an AIDS patient with a 2-year history of intestinal microsporidiosis. *J Clin Microbiol.* 1997; 35: 1862–1866. PMID: 9196210
7. Huibers MHW, Moons P, Maseko N, Gushu MB, Iwajomo OH, Heyderman RS, et al. Multiplex real-time PCR detection of intestinal protozoa in HIV-infected children in Malawi: *Enterocytozoon bieneusi* is common and associated with gastrointestinal complaints and may delay BMI (nutritional status) recovery. *Pediatr Infect Dis J.* 2018; 37: 910–915. <https://doi.org/10.1097/INF.0000000000001924> PMID: 29762365
8. Rogawski ET, Liu J, Platts-Mills JA, Kabir F, Lertsethtakarn P, Siguas M, et al. Use of quantitative molecular diagnostic methods to investigate the effect of enteropathogen infections on linear growth in children in low-resource settings: longitudinal analysis of results from the MAL-ED cohort study. *Lancet Glob Health.* 2018; 6:e1319–e1328. [https://doi.org/10.1016/S2214-109X\(18\)30351-6](https://doi.org/10.1016/S2214-109X(18)30351-6) PMID: 30287125
9. Santin M, Fayer R. Microsporidiosis: *Enterocytozoon bieneusi* in domesticated and wild animals. *Res Vet Sci.* 2011; 90: 363–371. <https://doi.org/10.1016/j.rvsc.2010.07.014> PMID: 20699192
10. Li W, Feng Y, Santin M. Host specificity of *Enterocytozoon bieneusi* and public health implications. *Trends Parasitol.* 2019; 35: 436–451. <https://doi.org/10.1016/j.pt.2019.04.004> PMID: 31076351

11. Li W, Feng Y, Zhang L, Xiao L. Potential impacts of host specificity on zoonotic or interspecies transmission of *Enterocytozoon bieneusi*. *Infect Genet Evol.* 2019;104033. <https://doi.org/10.1016/j.meegid.2019.104033> PMID: 31494271
12. Mathis A, Weber R, Deplazes P. Zoonotic potential of the microsporidia. *Clin Microbiol Rev.* 2005; 18: 423–445. <https://doi.org/10.1128/CMR.18.3.423-445.2005> PMID: 16020683
13. Izquierdo F, Castro Hermida JA, Fenoy S, Mezo M, González-Warleta M, del Aguila C. Detection of microsporidia in drinking water, wastewater and recreational rivers. *Water Res.* 2011; 45: 4837–4843. <https://doi.org/10.1016/j.watres.2011.06.033> PMID: 21774958
14. Santin M, Fayer R. *Enterocytozoon bieneusi* genotype nomenclature based on the internal transcribed spacer sequence: a consensus. *J Eukaryot Microbiol.* 2009; 56: 34–38. <https://doi.org/10.1111/j.1550-7408.2008.00380.x> PMID: 19335772
15. Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, Farag TH, Panchalingam S, et al. Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multi-center Study, GEMS): a prospective, case-control study. *Lancet.* 2013; 382: 209–222. [https://doi.org/10.1016/S0140-6736\(13\)60844-2](https://doi.org/10.1016/S0140-6736(13)60844-2) PMID: 23680352
16. Nkinin SW, Asonganyi T, Didier ES, Kaneshiro ES. Microsporidian infection is prevalent in healthy people in Cameroon. *J Clin Microbiol.* 2007; 45: 2841–2846. <https://doi.org/10.1128/JCM.00328-07> PMID: 17609328
17. Ikeh EI, Obadofin MO, Brindeiro B, Baugherb C, Frost F, Vanderjagt D, et al. Intestinal parasitism in Magama Gumau rural village and Jos township in north central Nigeria. *Niger Postgrad Med J.* 2007; 14: 290–295. PMID: 18163136
18. Saco C, Nhacolo A, Nhalungo D, Aponte JJ, Bassat Q, Augusto O, et al., Profile: Manhiça Health Research Centre (Manhiça HDSS). *Int J Epidemiol.* 2013; 42: 1309–1318. <https://doi.org/10.1093/ije/dyt148> PMID: 24159076
19. Alonso P, Saúte F, Aponte J, Gómez-Olivé F, Nhacolo A, Thomson R, et al. Manhiça DSS, Mozambique. Vol. 1, Population and health in developing countries. 2002. 189–195 p.
20. Kotloff KL, Blackwelder WC, Nasrin D, Nataro JP, Farag TH, van Eijk A, et al. The Global Enteric Multi-center Study (GEMS) of diarrheal disease in infants and young children in developing countries: epidemiologic and clinical methods of the case/control study. *Clin Infect Dis.* 2012; 55: S232–S245. <https://doi.org/10.1093/cid/cis753> PMID: 23169936
21. Panchalingam S, Antonio M, Hossain A, Mandomando I, Ochieng B, Oundo J, et al. Diagnostic microbiologic methods in the GEMS-1 case/control study. *Clin Infect Dis.* 2012; 55: S225–S231. <https://doi.org/10.1093/cid/cis787>
22. Buckholt MA, Lee JH, Tzipori S. Prevalence of *Enterocytozoon bieneusi* in swine: an 18-month survey at a slaughterhouse in Massachusetts. *Appl Environ Microbiol.* 2002; 68: 2595–2599. <https://doi.org/10.1128/aem.68.5.2595-2599.2002> PMID: 11976142
23. Kumar S., Stecher G., Li M., Niyaz C., Tamura K., Battistuzzi F.U. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol. Biol. Evol.* 2018; 35: 1547–1549. <https://doi.org/10.1093/molbev/msy096> PMID: 29722887
24. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol.* 1987; 4: 406–425. <https://doi.org/10.1093/oxfordjournals.molbev.a040454> PMID: 3447015
25. Heinze G, Schemper M. A solution to the problem of separation in logistic regression. *Stat Med.* 2002; 21:2409–2419. <https://doi.org/10.1002/sim.1047> PMID: 12210625
26. Irisarri-Gutiérrez MJ, Mingo MH, de Lucio A, Gil H, Morales L, Seguí R, et al. Association between enteric protozoan parasites and gastrointestinal illness among HIV- and tuberculosis-infected individuals in the Chowke district, southern Mozambique. *Acta Trop.* 2017; 170: 197–203. <https://doi.org/10.1016/j.actatropica.2017.03.010> PMID: 28302528
27. Meurs L, Polderman AM, Vinkeles Melchers NV, Brienen EA, Verweij JJ, Groosjohan B, et al. Diagnosing polyparasitism in a high-prevalence setting in Beira, Mozambique: Detection of intestinal parasites in fecal samples by microscopy and real-time PCR. *PLoS Negl Trop Dis.* 2017; 11: e0005310. <https://doi.org/10.1371/journal.pntd.0005310> PMID: 28114314
28. Augusto G, Nalá R, Casmo V, Sabonete A, Mapaco L, Monteiro J. Geographic distribution and prevalence of schistosomiasis and soil-transmitted helminths among schoolchildren in Mozambique. *Am J Trop Med Hyg.* 2009; 81: 799–803. <https://doi.org/10.4269/ajtmh.2009.08-0344> PMID: 19861614
29. Kotloff KL, Nasrin D, Blackwelder WC, Wu Y, Farag T, Panchalingam S, et al. The incidence, aetiology, and adverse clinical consequences of less severe diarrhoeal episodes among infants and children residing in low-income and middle-income countries: a 12-month case-control study as a follow-on to the Global Enteric Multicenter Study (GEMS). *Lancet Glob Health.* 2019; 7(5):e568–e584. [https://doi.org/10.1016/S2214-109X\(19\)30076-2](https://doi.org/10.1016/S2214-109X(19)30076-2) PMID: 31000128

30. Lobo ML, Augusto J, Antunes F, Ceita J, Xiao L, Codices V, et al. *Cryptosporidium* spp., *Giardia duodenalis*, *Enterocytozoon bieneusi* and other intestinal parasites in young children in Lobata province, Democratic Republic of São Tomé and Príncipe. PLoS One. 2014; 9: e97708. <https://doi.org/10.1371/journal.pone.0097708> PMID: 24846205
31. Ayinmode AB, Ojuromi OT, Xiao L. Molecular Identification of *Enterocytozoon bieneusi* isolates from Nigerian children. J Parasitol Res. 2011;129542. <https://doi.org/10.1155/2011/129542> PMID: 22132304
32. Samie A, Obi CL, Tzipori S, Weiss LM, Guerrant RL. Microsporidiosis in South Africa: PCR detection in stool samples of HIV-positive and HIV-negative individuals and school children in Vhembe district, Limpopo Province. Trans R Soc Trop Med Hyg. 2007; 101:547–554. <https://doi.org/10.1016/j.trstmh.2007.02.005> PMID: 17412378
33. Tumwine JK, Kekitiinwa A, Bakeera-Kitaka S, Ndeezi G, Downing R, Feng X, et al. Cryptosporidiosis and microsporidiosis in Ugandan children with persistent diarrhea with and without concurrent infection with the human immunodeficiency virus. Am J Trop Med Hyg. 2005; 73: 921–925. PMID: 16282304
34. Cegielski JP, Ortega YR, McKee S, Madden JF, Gaido L, Schwartz DA, et al. *Cryptosporidium*, *Enterocytozoon*, and *Cyclospora* infections in pediatric and adult patients with diarrhea in Tanzania. Clin Infect Dis. 1999; 28: 314–321. <https://doi.org/10.1086/515131> PMID: 10064250
35. ten Hove RJ, Van Lieshout L, Beadsworth MB, Perez MA, Spee K, Claas EC, et al. Characterization of genotypes of *Enterocytozoon bieneusi* in immunosuppressed and immunocompetent patient groups. J Eukaryot Microbiol. 2009; 56: 388–393 <https://doi.org/10.1111/j.1550-7408.2009.00393.x> PMID: 19602086
36. Breton J, Bart-Delabesse E, Biligui S, Carbone A, Seiller X, Okome-Nkoumou M, et al. New highly divergent rRNA sequence among biodiverse genotypes of *Enterocytozoon bieneusi* strains isolated from humans in Gabon and Cameroon. J Clin Microbiol. 2007; 45: 2580–2589. <https://doi.org/10.1128/JCM.02554-06> PMID: 17537939
37. Wang L, Zhang H, Zhao X, Zhang L, Zhang G, Guo M, et al. Zoonotic *Cryptosporidium* species and *Enterocytozoon bieneusi* genotypes in HIV-positive patients on antiretroviral therapy. J Clin Microbiol. 2013; 51: 557–563. <https://doi.org/10.1128/JCM.02758-12> PMID: 23224097
38. Sulaiman IM, Bern C, Gilman R, Cama V, Kawai V, Vargas D, et al. A molecular biologic study of *Enterocytozoon bieneusi* in HIV-infected patients in Lima, Peru. J Eukaryot Microbiol. 2003; 50: S591–S596.
39. Sarfati C, Bourgeois A, Menotti J, Liegeois F, Moyou-Somo R, Delaporte E, et al. Prevalence of intestinal parasites including microsporidia in human immunodeficiency virus-infected adults in Cameroon: a cross-sectional study. Am J Trop Med Hyg. 2006; 74:162–164. PMID: 16407362
40. Tumwine JK, Kekitiinwa A, Nabukeera N, Akiyoshi DE, Buckholt MA, Tzipori S. *Enterocytozoon bieneusi* among children with diarrhea attending Mulago Hospital in Uganda. Am J Trop Med Hyg. 2002; 67: 299–303. <https://doi.org/10.4269/ajtmh.2002.67.299> PMID: 12408671
41. Espern A, Morio F, Miegerville M, Illa H, Abdoulaye M, Meyssonier V, et al. Molecular study of microsporidiosis due to *Enterocytozoon bieneusi* and *Encephalitozoon intestinalis* among human immunodeficiency virus-infected patients from two geographical areas: Niamey, Niger, and Hanoi, Vietnam. J Clin Microbiol. 2007; 45: 2999–3002. <https://doi.org/10.1128/JCM.00684-07> PMID: 17634305
42. Maikai BV, Umoh JU, Lawal IA, Kudi AC, Ejembi CL, Xiao L. Molecular characterizations of *Cryptosporidium*, *Giardia*, and *Enterocytozoon* in humans in Kaduna State, Nigeria. Exp Parasitol. 2012; 131: 452–456. <https://doi.org/10.1016/j.exppara.2012.05.011> PMID: 22664352
43. Akinbo FO, Okaka CE, Omoregie R, Dearen T, Leon ET, Xiao L. Molecular epidemiologic characterization of *Enterocytozoon bieneusi* in HIV-infected persons in Benin City, Nigeria. Am J Trop Med Hyg. 2012; 86: 441–445. <https://doi.org/10.4269/ajtmh.2012.11-0548> PMID: 22403314
44. Ye J, Xiao L, Ma J, Guo M, Liu L, Feng Y. Anthroponotic enteric parasites in monkeys in public park, China. Emerg Infect Dis. 2012; 18: 1640–1643. <https://doi.org/10.3201/eid1810.120653> PMID: 23017776
45. Ben Ayed L, Yang W, Widmer G, Cama V, Ortega Y, Xiao L. Survey and genetic characterization of wastewater in Tunisia for *Cryptosporidium* spp., *Giardia duodenalis*, *Enterocytozoon bieneusi*, *Cyclospora cayetanensis* and *Eimeria* spp. J Water Health. 2012; 10: 431–444. <https://doi.org/10.2166/wh.2012.204> PMID: 22960487
46. Graczyk TK, Lucy FE, Mashinsky Y, Andrew Thompson RC, Koru O, Dasilva AJ. Human zoonotic enteropathogens in a constructed free-surface flow wetland. Parasitol Res. 2009; 105: 423–428. <https://doi.org/10.1007/s00436-009-1400-6> PMID: 19343366
47. Li N, Xiao L, Wang L, Zhao S, Zhao X, Duan L, et al. Molecular surveillance of *Cryptosporidium* spp., *Giardia duodenalis*, and *Enterocytozoon bieneusi* by genotyping and subtyping parasites in wastewater. PLoS Negl Trop Dis. 2012; 6: e1809. <https://doi.org/10.1371/journal.pntd.0001809> PMID: 22970334

48. Li W, Kiulia NM, Mwenda JM, Nyachio A, Taylor MB, Zhang X, et al. *Cyclospora papionis*, *Cryptosporidium hominis*, and human-pathogenic *Enterocytozoon bieneusi* in captive baboons in Kenya. *J Clin Microbiol*. 2011; 49: 4326–4329. <https://doi.org/10.1128/JCM.05051-11> PMID: 21956988
49. Widmer G, Dilo J, Tumwine JK, Tzipori S, Akiyoshi DE. Frequent occurrence of mixed *Enterocytozoon bieneusi* infections in humans. *Appl Environ Microbiol*. 2013; 79: 5357–5362. <https://doi.org/10.1128/AEM.01260-13> PMID: 23811516
50. Wang L, Xiao L, Duan L, Ye J, Guo Y, Guo M, et al. Concurrent infections of *Giardia duodenalis*, *Enterocytozoon bieneusi*, and *Clostridium difficile* in children during a cryptosporidiosis outbreak in a pediatric hospital in China. *PLoS Negl Trop Dis*. 2013; 7: e2437. <https://doi.org/10.1371/journal.pntd.0002437> PMID: 24069491
51. Ndzi ES, Asonganyi T, Nkinin MB, Xiao L, Didier ES, Bowers LC, et al. Fast technology analysis enables identification of species and genotypes of latent Microsporidia infections in healthy native Cameroonians. *J Eukaryot Microbiol*. 2016; 63: 146–152. <https://doi.org/10.1111/jeu.12262> PMID: 26303263
52. Raccurt CP, Agnamey P, Sarfati C, Chouaki T, Totet A. [Intestinal microsporidiosis due to *Enterocytozoon bieneusi* in a patient infected with HIV-1: first reported case in Chad]. *Med Trop*. 2008; 68: 307–308.
53. Wumba R, Jean M, Benjamin LM, Madone M, Fabien K, Josué Z, et al. *Enterocytozoon bieneusi* identification using real-time polymerase chain reaction and restriction fragment length polymorphism in HIV-infected humans from Kinshasa Province of the Democratic Republic of Congo. *J Parasitol Res*. 2012;278028. <https://doi.org/10.1155/2012/278028> PMID: 22811884
54. Wumba R, Longo-Mbenza B, Mandina M, Odio WT, Biligui S, Sala J, et al. Intestinal parasites infections in hospitalized AIDS patients in Kinshasa, Democratic Republic of Congo. *Parasite*. 2010; 17: 321–328. <https://doi.org/10.1051/parasite/2010174321> PMID: 21275238
55. Wumba R, Enache-Angoulvant A, Develoux M, Mulumba A, Mulumba PM, Hennequin C, et al. [Prevalence of opportunistic digestive parasitic infections in Kinshasa, Democratic Republic of Congo. Results of a preliminary study in 50 AIDS patients]. *Med Trop (Mars)*. 2007; 67: 145–148. PMID: 17691432
56. Endeshaw T, Kebede A, Verweij JJ, Wolday D, Zewide A, Tsige K, et al. Detection of intestinal microsporidiosis in diarrhoeal patients infected with the human immunodeficiency virus (HIV-1) using PCR and Uvitex-2B stain. *Ethiop Med J*. 2005; 43: 97–101. PMID: 16370539
57. Endeshaw T, Kebede A, Verweij JJ, Zewide A, Tsige K, Abraham Y, et al. Intestinal microsporidiosis in diarrheal patients infected with human immunodeficiency virus-1 in Addis Ababa, Ethiopia. *Jpn J Infect Dis*. 2006; 59: 306–310. PMID: 17060696
58. Mwachari C, Batchelor BI, Paul J, Waiyaki PG, Gilks CF. Chronic diarrhoea among HIV-infected adult patients in Nairobi, Kenya. *J Infect*. 1998; 37: 48–53. [https://doi.org/10.1016/s0163-4453\(98\)90561-8](https://doi.org/10.1016/s0163-4453(98)90561-8) PMID: 9733379
59. Greigert V, Abou-Bacar A, Brunet J, Nourrisson C, Pfaff AW, Benarbia L, et al. Human intestinal parasites in Mahajanga, Madagascar: The kingdom of the protozoa. *PLoS One* 2018; 13: e0204576. <https://doi.org/10.1371/journal.pone.0204576> PMID: 30304028
60. Maiga I, Doumbo O, Dembele M, Traoré H, Desportes-Livage I, Hilmarsdottir I, et al. [Human intestinal microsporidiosis in Bamako (Mali): the presence of *Enterocytozoon bieneusi* in HIV seropositive patients]. *Sante*. 1997; 7: 257–262. PMID: 9410452
61. Alfa Cisse O, Ouattara A, Thellier M, Accoceberry I, Biligui S, Minta D, et al. Evaluation of an immunofluorescent-antibody test using monoclonal antibodies directed against *Enterocytozoon bieneusi* and *Encephalitozoon intestinalis* for diagnosis of intestinal microsporidiosis in Bamako (Mali). *J Clin Microbiol*. 2002; 40: 1715–1718. <https://doi.org/10.1128/jcm.40.5.1715-1718.2002> PMID: 11980948
62. Bretagne S, Foulet F, Alkassoum W, Fleury-Feith J, Develoux M. [Prevalence of *Enterocytozoon bieneusi* spores in the stool of AIDS patients and African children not infected by HIV]. *Bull Soc Pathol Exot*. 1993; 86: 351–357. PMID: 8124104
63. Ayinmode AB, Zhang H, Dada-Adegbola HO, Xiao L. *Cryptosporidium hominis* subtypes and *Enterocytozoon bieneusi* genotypes in HIV-infected persons in Ibadan, Nigeria. *Zoonoses Public Health*. 2014; 61: 297–303. <https://doi.org/10.1111/zph.12072> PMID: 23870732
64. Akinbo FO, Okaka CE, Omoregie R, Adamu H, Xiao L. Unusual *Enterocytozoon bieneusi* genotypes and *Cryptosporidium hominis* subtypes in HIV-infected patients on highly active antiretroviral therapy. *Am J Trop Med Hyg*. 2013; 89: 157–161. <https://doi.org/10.4269/ajtmh.12-0635> PMID: 23629938
65. Ojuromi OT, Izquierdo F, Fenoy S, Fagbenro-Beyioku A, Oyibo W, Akanmu A, et al. Identification and characterization of microsporidia from fecal samples of HIV-positive patients from Lagos, Nigeria. *PLoS One*. 2012; 7: e35239. <https://doi.org/10.1371/journal.pone.0035239> PMID: 22496910

66. Ojuromi OT, Duan L, Izquierdo F, Fenoy SM, Oyibo WA, Del Aguila C, et al. Genotypes of *Cryptosporidium* spp. and *Enterocytozoon bieneusi* in human immunodeficiency virus-infected patients in Lagos, Nigeria. *J Eukaryot Microbiol*. 2016; 63: 414–418. <https://doi.org/10.1111/jeu.12285> PMID: 26662459
67. Chabchoub N, Abdelmalek R, Mellouli F, Kanoun F, Thellier M, Bouratbine A, et al. Genetic identification of intestinal microsporidia species in immunocompromised patients in Tunisia. *Am J Trop Med Hyg*. 2009; 80: 24–27. PMID: 19141834
68. Abdelmalek R, Anane S, Chabchoub N, Essid R, Aoun K, Chaabéne TB, et al. [Microsporidia and cryptosporidia coinfection in an HIV-infected newborn]. *Arch Pediatr*. 2011; 18: 562–564. <https://doi.org/10.1016/j.arcped.2011.02.016> PMID: 21458971
69. Anane S, Kaouech E, Belhadj S, Abdelmalek R, Ammari L, Ben Othman T, et al. [Identification of *Enterocytozoon bieneusi* by PCR in stools of Tunisian immunocompromised patients]. *Pathol Biol*. 2011; 59: 234–239. <https://doi.org/10.1016/j.patbio.2009.03.007> PMID: 19481368
70. Chabchoub N, Abdelmalek R, Issa S, Kanoun F, Ben Chaabene T, Bouratbine A, et al. Apport de la PCR dans la recherche et l'identification des microsporidies intestinales chez les sujets infectés par le VIH. *Pathol Biol*. 2012; 60: 91–94. <https://doi.org/10.1016/j.patbio.2009.07.034>
71. Chabchoub N, Abdelmalek R, Breton J, Kanoun F, Thellier M, Bouratbine A, et al. Genotype identification of *Enterocytozoon bieneusi* isolates from stool samples of HIV-infected Tunisian patients. *Parasite*. 2012; 19: 147–151. <https://doi.org/10.1051/parasite/2012192147> PMID: 22550625
72. Aissa S, Chabchoub N, Abdelmalek R, Kanoun F, Goubantini A, Ammari L, et al. Asymptomatic intestinal carriage of microsporidia in HIV-positive patients in Tunisia: prevalence, species, and pathogenesis. *Med Sante Trop*. 2017; 27: 281–285. <https://doi.org/10.1684/mst.2017.0697> PMID: 28947404
73. Lucas SB, Papadaki L, Conlon C, Sewankambo N, Goodgame R, Serwadda D. Diagnosis of intestinal microsporidiosis in patients with AIDS. *J Clin Pathol*. 1989; 42: 885–887.
74. Hautvast JL, Tolboom JJ, Derks TJ, Beckers P, Sauerwein RW. Asymptomatic intestinal microsporidiosis in a human immunodeficiency virus-seronegative, immunocompetent Zambian child. *Pediatr Infect Dis J*. 1997; 16: 415–416. <https://doi.org/10.1097/00006454-199704000-00017> PMID: 9109148
75. van Gool T, Luderhoff E, Nathoo KJ, Kiire CF, Dankert J, Mason PR. High prevalence of *Enterocytozoon bieneusi* infections among HIV-positive individuals with persistent diarrhoea in Harare, Zimbabwe. *Trans R Soc Trop Med Hyg*. 1995; 89: 478–480. [https://doi.org/10.1016/0035-9203\(95\)90073-x](https://doi.org/10.1016/0035-9203(95)90073-x) PMID: 8560514
76. Carville A, Mansfield K, Widmer G, Lackner A, Kotler D, Wiest P, et al. Development and application of genetic probes for detection of *Enterocytozoon bieneusi* in formalin-fixed stools and in intestinal biopsy specimens from infected patients. *Clin Diagn Lab Immunol*. 1997; 4: 405–408. PMID: 9220155
77. Gumbo T, Sarbah S, Gangaidzo IT, Ortega Y, Sterling CR, Carville A, et al. Intestinal parasites in patients with diarrhea and human immunodeficiency virus infection in Zimbabwe. *AIDS*. 1999; 13: 819–821. <https://doi.org/10.1097/00002030-199905070-00011> PMID: 10357381
78. Gumbo T, Gangaidzo IT, Sarbah S, Carville A, Tzipori S, Wiest PM. *Enterocytozoon bieneusi* infection in patients without evidence of immunosuppression: two cases from Zimbabwe found to have positive stools by PCR. *Ann Trop Med Parasitol*. 2000; 94: 699–702. <https://doi.org/10.1080/00034983.2000.11813593> PMID: 11144811